

# Convexity check

February 1, 2006

## 1 Preliminaries

Recall Expression (22) in our current paper:

$$\frac{\partial \ln F_m}{\partial m_{i,r}^{j,t}} = \frac{1}{2} \ln m_{i,r}^{j,t} - \ln \mu_{i,r}^{j,t} + \frac{3}{10} \quad (1)$$

Taking advantage:

$$m_{i,r}^{j,t} = \mu_{i,r}^{j,t} \frac{\partial \Phi(\tilde{\mu})}{\partial \mu_{i,r}^{j,t}} \Leftrightarrow m_{i,r}^{j,t} = \frac{\partial \Phi(\tilde{\mu})}{\partial \ln \mu_{i,r}^{j,t}} \quad (2)$$

we can get:

$$\nabla_{m_{i,r}^{j,t}}^2 \ln F_m = \left[ \frac{1}{2m_{i,-1}^{j,-1}}, \frac{1}{2m_{i,-1}^{j,1}}, \frac{1}{2m_{i,1}^{j,-1}}, \frac{1}{2m_{i,1}^{j,1}} \right] \times \mathbb{I}_{4,4} - \left( \nabla_{\ln \mu_{i,r}^{j,t}}^2 \Phi(\tilde{\mu}) \right)^{-1} \quad (3)$$

We wish to show that the LPDMs of (3) alternate properly in sing, for each 4tuple of  $m > 0$ 's (which implicitly determines the 4tuple of  $\mu$ 's) that sum to a fixed positive constant  $n_i^j > 0$ , i.e. under the constraints:

$$\begin{aligned} m_{i,r}^{j,t} + m_{i,r}^{j,-t} &\leq 4 \left( m_{i,-r}^{j,t} + m_{i,-r}^{j,-t} \right) \\ m_{i,r}^{j,t} + m_{i,-r}^{j,t} &\leq 4 \left( m_{i,-r}^{j,-t} + m_{i,-r}^{j,t} \right) \\ m_{i,r}^{j,t} &= m_{i+r,-r}^{j+t,-t} \\ \sum_{r,t} m_{i,r}^{j,t} &= n_i^j \end{aligned} \quad (4)$$

A nice observation is that we can take advantage of (2) and transform (3) to (5), as a function of  $\mu$ 's only. Then proper alternation of signs of (5), for *all positive*  $\mu$ 's, will imply that the LPDMs of (3) alternate properly in sing for each 4tuple of  $m$ 's that satisfy constraints (4).

$$\nabla_{m_{i,r}^{j,t}}^2 \ln F_m = \left[ \frac{1}{2\mu_{i,-1}^{j,-1} \frac{\partial \Phi(\tilde{\mu})}{\partial \mu_{i,-1}^{j,-1}}}, \frac{1}{2\mu_{i,-1}^{j,1} \frac{\partial \Phi(\tilde{\mu})}{\partial \mu_{i,-1}^{j,1}}}, \frac{1}{2\mu_{i,1}^{j,-1} \frac{\partial \Phi(\tilde{\mu})}{\partial \mu_{i,1}^{j,-1}}}, \frac{1}{2\mu_{i,1}^{j,1} \frac{\partial \Phi(\tilde{\mu})}{\partial \mu_{i,1}^{j,1}}} \right] \times \mathbb{I}_{4,4} - \left( \nabla_{\ln \mu_{i,r}^{j,t}}^2 \Phi(\tilde{\mu}) \right)^{-1} \quad (5)$$

Observe that each entry in (5) is a *regular* polynomial, i.e. it consists of monomials that their exponents sum to a fixed constant. Then to show proper alternation of signs of (5) for *all positive*  $\mu$ 's, it suffices to show it

for all 4tuples of  $\mu$ 's that sum *only* to 1. For example, let:  $f(x, y, z, w) = x^2y^2zw^3 - 10xyz^3w^5 + x^5w^3$  with exponents summing to 8. Then,

$$\begin{aligned} f(x, y, z, w) &= (x^2y^2zw^3 - 10xyz^3w^5 + x^5w^3) \frac{(x + y + z + w)^8}{(x + y + z + w)^8} \\ &= (p_x^2p_y^2p_zp_w^3 - 10p_xp_y p_z^3p_w^5 + p_x^5p_w^3) (x + y + z + w)^8 \\ &\text{with: } (x + y + z + w)^8 > 0 \end{aligned} \tag{6}$$

where  $p_t = \frac{t}{x+y+z+w} \in (0, 1]$  for each subscript  $t \in \{x, y, z, w\}$ . In other words, the sign of (6) depends *only* on the scaled vector  $(p_x, \dots, p_w) \in (0, 1]^4$  of the positive variables  $x, y, z, w$ .

**The C program:** We represent the 4  $\mu$ 's as  $\mu_1, \dots, \mu_4$  and the 4  $m$ 's as  $m_1, \dots, m_4$ . Then  $\mu_1$  ranges in  $[bound, 1 - bound]$  with **step**= 1/5000 and **bound**= 0.00000001. For each fixed value of  $\mu_1$  then  $\mu_2$  ranges in  $[bound, 1 - \mu_1]$  with **step**= 1/5000. For each fixed tuple of values  $(\mu_1, \mu_2)$  then  $\mu_3$  ranges in  $[bound, 1 - \mu_1 - \mu_2]$  with **step**= 1/5000. Finally, for each tuple of values  $(\mu_1, \mu_2, \mu_3)$  then we set  $\mu_4 = 1 - \mu_1 - \mu_2 - \mu_3 \geq bound$ .

The 2 matrices in Expression (5) are implemented as follows. The leftmost matrix

$$\left[ \begin{array}{cccc} 1 & 1 & 1 & 1 \\ 2\mu_1 \frac{\partial \Phi(\tilde{\mu})}{\partial \mu_1} & 2\mu_2 \frac{\partial \Phi(\tilde{\mu})}{\partial \mu_2} & 2\mu_3 \frac{\partial \Phi(\tilde{\mu})}{\partial \mu_3} & 2\mu_4 \frac{\partial \Phi(\tilde{\mu})}{\partial \mu_4} \end{array} \right] \times \mathbf{I}_{4,4}$$

in (5) is evaluated per tuple of  $m\mu$ 's in the C program in the  $4 \times 4$  matrix **Minvf**. The rightmost matrix

$$\nabla_{\ln \mu_1, \dots, \ln \mu_4}^2 \Phi(\tilde{\mu})$$

in (5) is evaluated per tuple of  $m\mu$ 's in the C program as the  $4 \times 4$  matrix **Sf**, while its corresponding inverse in  $4 \times 4$  matrix **Sinvf**. Since  $\mu$ 's add to 1, it is convenient to use the  $3 \times 3$  matrix

$$HG = A \times (\mathbf{Minvf} - \mathbf{Sinf}) \times \text{transpose}(A)$$

where:

$$A = \begin{bmatrix} 1 & 0 & 0 & -1 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & -1 \end{bmatrix}$$

Finally, the LPDMs of matrix **HG** are denoted as **currentLPDM1**,  $\dots$ , **currentLPDM3**.